

PHOTOMETRIC ANALYSIS AND ROTATION PERIOD DETERMINATION FOR ASTEROIDS 5445 WILLIWAW, (8823) 1987 WS3 AND (26568) 2000 ET49

Alessandro Marchini
Astronomical Observatory, DSFTA - University of Siena (K54)
Via Roma 56, 53100 - Siena, ITALY
marchini@unisi.it

Riccardo Papini
Wild Boar Remote Observatory (K49)
San Casciano in Val di Pesa (FI), ITALY

Giulio Scarfi
Iota Scorpil Observatory (K78), La Spezia, ITALY

(Received: 2021 Jan 15)

Photometric observations of three main-belt asteroids were conducted in order to determine their synodic rotation periods. For 5445 Williwaw we found $P = 10.65 \pm 0.02$ h, $A = 0.29 \pm 0.05$ mag; for (8823) 1987 WS3, a slow rotator, we found a rough period of $P = 86.0 \pm 1.0$ h, $A = 0.24 \pm 0.02$ mag; for (26568) 2000 ET49 we found $P = 7.171 \pm 0.003$ h, $A = 0.71 \pm 0.04$ mag



CCD photometric observations of three main-belt asteroids were carried out in 2020 October - December at three Italian observatories. At the Astronomical Observatory of the University of Siena (K54), a facility inside the Department of Physical Sciences, Earth and Environment (DSFTA, 2020), we used a 0.30-m $f/5.6$ Maksutov-Cassegrain telescope, SBIG STL-6303E NABG CCD camera, and clear filter; the pixel scale was 2.30 arcsec when binned at 2×2 pixels and all exposures were 300 seconds. At the Wild Boar Remote Observatory (K49) data were obtained with a 0.235-m $f/10$ (SCT) telescope, a SBIG ST8-XME NABG CCD camera unfiltered; the pixel scale was 1.60 arcsec in binning 2×2 and all exposures were 300 seconds. At the Iota Scorpil Observatory (K78) we used a 0.40-m $f/6$ Ritchey-Chretien telescope, SBIG STXL 6303E NABG CCD camera, and R filter; the pixel scale was 1.55 when binned 2×2 and all exposures were 180 seconds.

Data processing and analysis were done with *MPO Canopus* (Warner, 2018). All images were calibrated with dark and flat-field frames and the instrumental magnitudes converted to R magnitudes using solar-colored field stars from a version of the CMC-15 catalogue distributed with *MPO Canopus*. Table I shows the observing circumstances and results.

A search through the asteroid lightcurve database (LCDB; Warner et al., 2009) indicates that our results may be the first reported lightcurve observations and results for these asteroids.

5445 Williwaw. (1991 PA12) was discovered on 1991 August 7 by H.E. Holt at Mount Palomar and named after a dramatic mountain on the skyline of Anchorage. Mount Williwaw stands 5445 feet above sea level and it's the highest point in the Campbell Creek drainage. [Ref: Minor Planet Circ. 34341] It is a main-belt asteroid with a semi-major axis of 2.552 AU, eccentricity 0.223, inclination 6.115° , and an orbital period of 4.08 years. Its absolute magnitude is $H = 12.4$ (JPL, 2020). The WISE/NEOWISE satellite infrared radiometry survey (Masiero et al., 2014) found a diameter $D = 8.797 \pm 0.107$ km using an absolute magnitude $H = 12.2$.

Observations were conducted over three nights and collected 213 data points. The period analysis shows a solution for the rotational period of $P = 10.65 \pm 0.02$ h with an amplitude $A = 0.29 \pm 0.05$ mag.




Number	Name	2020/mm/dd	Phase	L_{PAB}	B_{PAB}	Period(h)	P.E.	Amp	A.E.	Grp
5445	Williwaw	10/17-10/20	2.4, 2.6	24	6	10.65	0.02	0.29	0.05	MB
8823	1987 WS3	11/18-12/15	*6.8, 10.0	68	-2	86.0	1.0	0.24	0.02	MB
26568	2000 ET49	11/06-11/10	4.4, 5.3	41	-8	7.171	0.003	0.71	0.04	MB

Table I. Observing circumstances and results. The first line gives the results for the primary of a binary system. The second line gives the orbital period of the satellite and the maximum attenuation. The phase angle is given for the first and last date. If preceded by an asterisk, the phase angle reached an extrema during the period. L_{PAB} and B_{PAB} are the approximate phase angle bisector longitude/latitude at mid-date range (see Harris et al., 1984). Grp is the asteroid family/group (Warner et al., 2009).



(8823) 1987 WS₃. (1981 QC1) was discovered on 1987 November 24 by S. McDonald at Anderson Mesa. It is a main-belt asteroid with a semi-major axis of 2.570 AU, eccentricity 0.240, inclination 13.557°, and an orbital period of 4.12 years. Its absolute magnitude is $H = 12.8$ (JPL, 2020). The WISE/NEOWISE satellite infrared radiometry survey (Masiero et al., 2011) found a diameter $D = 10.652 \pm 0.108$ km using an absolute magnitude $H = 12.6$.

Observations over seven nights collected 351 data points. The period analysis shows a possible bimodal solution for the rotational period of $P = 86.0 \pm 1.0$ h with an amplitude $A = 0.24 \pm 0.02$ mag. This target revealed to be a very slow rotator and the result is based on less than full coverage, so that the true period may differ a few hours.



(26568) 2000 ET₄₉ was discovered on 2000 March 9 by LINEAR at Socorro. It is a main-belt asteroid with a semi-major axis of 3.155 AU, eccentricity 0.208, inclination 14.842°, and an orbital period of 5.60 years. Its absolute magnitude is $H = 13.0$ (JPL, 2020). The WISE/NEOWISE satellite infrared radiometry survey (Masiero et al., 2011) found a diameter $D = 16.708 \pm 0.160$ km using an absolute magnitude $H = 13.0$.

Observations were conducted over three nights and collected 245 data points. The period analysis shows a result for the rotational period of $P = 7.171 \pm 0.003$ h with an amplitude $A = 0.71 \pm 0.04$ mag as the most likely bimodal solution for this asteroid.



Acknowledgements

Minor Planet Circulars (MPCs) are published by the International Astronomical Union's Minor Planet Center.
https://www.minorplanetcenter.net/iau/ECS/MPCArchive/MPCArchive_TBL.html

References

- DSFTA (2020). Dipartimento di Scienze Fisiche, della Terra e dell'Ambiente - Astronomical Observatory.
<https://www.dsfta.unisi.it/en/research/labs/astronomical-observatory>
- Harris, A.W.; Young, J.W.; Scaltriti, F.; Zappala, V. (1984). "Lightcurves and phase relations of the asteroids 82 Alkmene and 444 Gyptis." *Icarus* **57**, 251-258.
- JPL (2020). Small-Body Database Browser.
<http://ssd.jpl.nasa.gov/sbdb.cgi#top>
- Masiero, J.R.; Mainzer, A.K.; Grav, T.; Bauer, J.M.; Cutri, R.M.; Dailey, J.; Eisenhardt, P.R.M.; McMillan, R.S.; Spahr, T.B.; Skrutskie, M.F.; Tholen, D.; Walker, R.G.; Wright, E.L.; DeBaun, E.; Elsbury, D.; Gautier IV, T.; Gomillion, S.; Wilkins, A. (2011). "Main Belt Asteroids with WISE/NEOWISE. I. Preliminary Albedos and Diameters." *Astrophys. J.* **741**, A68.